The logic of Comparative Theory Evaluation

by M.L.C. Redhead, Chelsea College, London.

25.10.75

Abstract

Schaffner's logic of comparative theory evaluation is criticised for its incorrect interpretation of Bayes's theorem. A corrected analysis is given and extended to the case of multiple successful predictions by a theory. The application of the method to the appraisal of quantitative theories is discussed. Finally two Bayesian models are presented for the general case in which successful and unsuccessful predictions are interspersed in the development of a research programme. An alternative approach incorporating the notion of methodological decision is discussed and simple representations are provided for the life and death of a research programme.

A Bayesian model is next presented for the general case in which successful and unsuccessful predictions are interspersed in the development of a research programme. An alternative approach incorporating the notion of methodological decision is discussed and simple representations are provided for the life and death of a research programme.

The logic of Comparative Theory Evaluation

1. A Critique of Schaffner's Account

In a recent note Schaffner ([1974]) has [discussed] given a formal discussion of the notion of ad hoc in terms of a Bayesian model for the appraisal of theories. Schaffner develops his general ideas in the context of a critique of Zahar's [1973] which was concerned with the particular problem of comparing the Einstein and Lorentz research programmes. Zahar suggests the following analysis of ad hocness:[1]

ad hoc: A theory is said to be ad hoc, if it has no novel consequences as compared with its predecessor

ad hoc$_2$: [A theory] ... is ad hoc$_2$ if none of its novel predictions have been actually 'verified'.

ad hoc$_3$: [A] ... theory is said to be ad hoc$_3$ if it is obtained from its predecessor through a modification of the auxiliary hypotheses which does not accord with the spirit of the heuristic of the programme.

Zahar explains the meaning of novelty as follows:[2]

A fact will be considered novel with respect to a given hypothesis if it did not belong to the problem-situation which governed the construction of the hypothesis.

Schaffner's attempted elucidation is really rather confused. He first discusses the notions of ad hoc$_1$ and ad hoc$_3$. The

[1] Zahar [1973] p. 101

[2] Zahar [1973] p. 103

first he describes as a logical dream, since the penal consequences of a theory cannot in [the] practice be "surveyed" so the question of whether a theory is ad hoc, can only be discussed related to the extent to which penal consequences have been looked for at the particular epoch of the evaluation. Although Schaffner claims to be "vague" to the point of inapplicability. For Schaffner ad hoc is "clear" to the sense in which ad hoc is used in science and he announces that his Bayesian apologies will be brought to bear on this sense of ad hoc.

He begins his analysis[3] by referring to the general division theorem by which he actually means Bayes's theorem. Schaffner says that this is a non-controversial theorem in the probability calculus and refers to Salmon's [1967] account.

He claims in a footnote[4] to have improved on Salmon's presentation, "unfortunately" so far "improves" the theorem that he actually renders it false. Denoting by $P(T/e\&b)$ the probability of a theory T to be true in the light of background knowledge b and the factual outcome e of some experiment get part of by $P(T/b)$ the prior assessment of T, by $P(e/T\&b)$ the probability of obtaining e given T and b, and by $P(e/b)$ "the probability of obtaining e on the basis of theories other than T", Schaffner writes

$$P(T/e\&b) = P(T/b) \cdot \frac{P(e/T\&b)}{P(e/b)}$$

(Schaffner's theorem)

Ibid

[3] Schaffner [1974] p. 69–73

[4] Footnote 3 p. 70.

(1) $p(T/a \& e) = \dfrac{p(T/a) \cdot p(e/T \& a)}{p(e/a)}$ (A)

In a footnote[5] he explains $p(e/a)$ can be expanded in terms of a pt of n "alternative competing theories" (my italics) in the form

(2) $p(e/a) = \sum_{i=1}^{n} p(T_i/a)\, p(e/T_i \& a)$.

Now

Now (1) is of course true only if $p(e/a)$ includes a contribution from itself, which Schaffer explicitly denies[6] in his definition of $p(e/a)$. This unfortunate oversight now makes nonsense of Schaffer's subsequent analysis. Because he assumes $p(e/a)$ is independent of $p(T/a)$ he makes his criterion for a theory to be ad hoc$_2$ (Footnote 6) Schaffer actually applies in terms of hypotheses being ad hoc but we want to reach the argument as simply as possible and depend essentially on the smallness of $p(T/a)$. The argument[a] for $p(T/a)$ being small is essentially that it has no "theoretical support". This looks superficially like a reference to ad hoc$_3$. And Schaffer now appears to be claiming[b] that a theory is ad hoc$_2$ [in virtue of its being] ad hoc$_3$. So[c] he is not really giving an independent analysis[d] of ad hoc$_2$ at all. But it is pointless to criticise his detailed arguments given[e] how all derived[f] by using

5 & 6 Quid Footnote 2 p. 70.

Insertion
new Ford

Note that we follow Schaffner in interpreting $p(A/B)$ as probable that A is true given that B is true.

Note also that e is subscripted on quotes (3) to refer to a prediction made by the theory T (as used before note e_T to emphasize this [deleted] important point) — $p(e(n\,T\,\&\,?))$ means [deleted] the probability that the predicted e_T deduced from the theory T is a true prediction given that the the background information b is true but the theory T is false (i.e. its consequences are not guaranteed to be true although "by accident" they may be true)

the wrong formulation of Bayes's theorem. ☐
The only statement which Scheffler sets right is where he discusses the "odds form" for comparing the evidential of a theory with its rivals*7. Here in taking the appropriate ratio p(e/h) cancels out thus is the one point at which her argument is not invalidated by her incorrect analysis of p(e/h).

2. The Bayesian Analysis of Adhocness

It is not just a fairly trifle matter to get an "internal" analysis of adhoc, they were the correct form of Bayes's theorem, which we write as Bayes's

(3) $$P(T/a\&e) = \frac{P(T/a)\cdot P(e/T\&a)}{P(e/T\&a)\,P(T/a) + P(e/\sim T\&a)\,P(\sim T/a)}$$

where $\sim T$ denotes the negation of T.

Putting $P(T/a) = x$, $P(e/\sim T\&a) = \varepsilon$

and taking $P(e/T\&a) = 1$ and using $P(\sim T/a) = 1 - x$ yields

(4) $$P(T/a\&e) = \frac{x}{x + \varepsilon(1-x)}$$

We define an enhancement rate Λ by

$$\Lambda = \frac{P(T/a\&e)}{P(T/a)}$$

where using (4)

*7. Ibid p 72.

we obtain the simple result

$$(5) \qquad \Omega = \frac{1}{x + \varepsilon(1-x)} .$$

new word
We can now explain that a theory T is ad hoc with respect to the experiment e if $\varepsilon = 1$, i.e. if the explanation of e by T in no way depends on the truth or falsehood of T but of what eventualities but with certainty to the result e.
This is just what a scientist means when he says T was an ad hoc explanation of e, namely T was devised for the express purpose of explaining e, so the explanation of e is guaranteed independent of whether T is true or false.
To show the consistency of our analysis if we put $\varepsilon = 1$ in (5) we get $\Omega = 1$ so the posterior and prior probabilities of T are equal (there is no enhancement) and this again is just what we expect from an ad hoc explanation of e namely e {~~gives us no~~} tells us no additional information for assessing the truth of T.

new word
We notice that if $x \leq \varepsilon \leq 1$, then $\Omega \geq 1/\varepsilon$ so in this case we get a big enhancement and the theory is

for from being ad hoc, this is just the situation in which Schaffner would claim that the theory was ad hoc, which highlights the way in which her analysis goes astray

3. The case of multiple Predictions

To develop our analysis a little further we can consider how a theory builds up a favourable appraisal as it makes a number of successful predictions $e_1, e_2 \ldots e_n$ say. Denote by P_s the posterior probability $P(T/e_1 \& e_2 \ldots \& e_s)$ after s successful predictions. Assuming the predictions are quite independent and [~~executed with~~] for simplicity are all associated with the same value of ε, we can clearly write

$$P_n = \mathcal{R}^{(n)} \times \mathcal{R}^{(n-1)} \times \ldots \mathcal{R}^{(1)} \cdot P_0$$

where $P_0 = x$ according to our previous notation

and
$$\left.\begin{aligned} \mathcal{R}^{(s)} &= \frac{1}{P_{s-1}(1-\varepsilon)+\varepsilon} \\ P_{s+1} &= \mathcal{R}^{(s+1)} \cdot P_s \end{aligned}\right\}$$

The solution of this recursion is by induction (or more simply by replacing $\varepsilon \to \varepsilon^n$ in the formula for P_1 (see (4) above). We obtain

$$(6) \qquad P_n = \frac{1}{1-\varepsilon^n+\varepsilon^n/x}$$

We can also ask what is the probability [~~obtaining~~] the $(n+1)^{th}$ prediction e_{n+1} being correct if the theory has already made n

successful predictions $e_1 \ldots e_n$. Denoting their probability by $P(e_{n+1})$ we easily obtain the result

$$P(e_{n+1}) = \varepsilon + \frac{1-\varepsilon}{1-\varepsilon^n + \varepsilon^n/x} \tag{6}$$

If ε is a small quantity ($i.e. \ll 1$) [deleted: which] would be the case if T is non-adiabatic with respect to all the predictions, and can create the following approximate formulae which would be perfectly satisfactory for the subsequent discussion

$$P_n \simeq \frac{1}{1+\varepsilon^n/x} \tag{7}$$

$$P(e_{n+1}) \simeq \varepsilon + P_n \tag{8}$$

$$\Omega^{(n)} \simeq \frac{1}{\varepsilon + P_{n-1}} \tag{9}$$

We note the following features

(i) The value of P_n depends critically on the ratio ε^n/x.

For $\varepsilon^n/x \gg 1$, $P_n \ll 1$

and for $\varepsilon^n/x \ll 1$, $P_n \simeq 1$

At the critical point $\varepsilon^n = x$, we have $P_n \simeq 1/2$.

So if initially $x \ll \varepsilon$ as n increases P_n will rise steeply as we reach the value $n = \ln x / \ln \varepsilon$.

(2) So long as $P_n \leq \varepsilon$, we have $P(e_{n+1}) \geq \varepsilon$, but as P_n builds up towards unity, so does $P(e_{n+1})$

(3) So long as $P_{n-1} \leq \varepsilon$, $\lambda^{(n)} \leq 1/\varepsilon$, but as P_{n-1} builds up towards unity the enhancement factor $\lambda^{(n)}$ also tends to unity

new para

To take a concrete ~~illustration~~ case we illustrate in Fig 1. P_n and $P(e_{n+1})$ as functions of n for the particular choice $x = .01$, $\varepsilon = -1$.



Fig 1.

4. Quantitative Predictions

We can apply this analysis to the important case of quantitative predictions.

Suppose a theory T predicts exactly an experimental result which is known to an accuracy of n significant figures. We assume as part of our background

knowledge is that the order of magnitude of the result is known, i.e. we disregard the prediction of zeros occurring before or after the significant figures in the experimental result.

As a concrete illustration we cite the theoretical predictions of the anomaly of the hydrogen spectrum known as the Lamb shift and the anomaly in the magnetic moment of the electron. For example the latter is now known experimentally to be 0.00115965 7.7 ± 3.5 Bohr magnetons whereas the theoretical prediction is 0.00115965 5.4 ± 3.3 magnetons. (For a good account of the fluctuating agreement between theory and experiment for the anomaly see reviews by Lautrup, Peterman and de Rafael in Physics Reports [1972] if you are interested.)

Clearly if we regard the prediction of each significant figure as an independent event then the difference value to take for ε is .1 since a false theory could have ten equal probabilities for being in each digit. The question of what value to take for x is somewhat arbitrary. We follow Schaffner in adopting a purely logical approach and set x = 0. For our purposes x reflects the scientific confidence in the new theory. One could argue that a scientist would not spend great efforts developing the consequences of a theory he did not believe in. By analogy with the skeptic

in the Bayesian analysis of significance testing (see for example Pollard [1974]) we could take $\alpha = 1/2$. Perhaps more realistically one should take α around 0.01 and adopt the sociological rule that values a scientist has a one percent level of confidence in the truth of his theory he would not seriously investigate it.

With this choice of parameters we see that the build up of confidence in a theory which makes exact quantitative predictions, as the accuracy of the experiment increases, would be characterized by the steepness for P_n in Fig. 1.

Of course $P(\theta_{n+1})$ is only given by the analysis so long as other factors which could potentially influence the result are known not to be significant. For a certain value of n this condition will fail. For example in the case of the anomalous magnetic moment of the electron the effect of hadronic corrections introduces theoretical uncertainties which would ultimately make the prediction of the theory unreliable.[8]

On the other hand the Lamb shift has so far been measured with an accuracy less by a factor of twenty, so the physicist's confidence in a refined quantitative prediction here is very high.

---

[8] According to Kiel, Verley [1972] the known hadronic contributions to the electron anomaly would affect the tenth significant figure.

5. A Bayesian Model for the Methodology of Research Programmes

We now want to extend our analysis to the case where the theory T consists of two parts

$T = C \,\&\, B_i$ where C is the hard core and B_i an auxiliary hypothesis selected from a set $\{B_i\}$ which represents the particle articulations & the protective belt in Lakatos's terminology (see for example Rao [1976])

We write

$$P(C/e) = \frac{P(e/c)\,P(c)}{P(e/c)\,P(c) + P(e/\neg c)\,P(\neg c)}$$

and expand $P(e/c) = P(e/c \,\&\, B)\,P(B/c) + P(e/c \,\&\, \neg B)\,P(\neg B/c)$

(Footnote 9. This method of analysing Bayes's theorem was suggested to me by Jon Dorling)

Introducing abbreviations $P(c) = x$, $P(B) = y$,

$P(e/c \,\&\, \neg B) = R_1$,

$P(e/\neg c) = R_2$

and setting $P(e/c \,\&\, B) = 1$, $P(\neg B) = 1-y$, $P(\neg c) = 1-x$

the enhancement ratio for hard core is λ_c

$$\text{where } \lambda_c = \frac{P(c/e)}{x} = \frac{y + R_1(1-y)}{\big(y + R_1(1-y)\big)x + R_2(1-x)}$$

If we take $R_1 = R_2 = R$ say we get

$$(10)\quad \lambda_c = \frac{y + R(1-y)}{\big(y + R(1-y)\big)x + R(1-x)} \;\leq\; \frac{y + R(1-y)}{xy + R(1-xy)}$$

(the enhancement ratio for the auxiliary hypotheses)

Similarly for the auxiliaries λ_B, in the form

$$(11)\quad \lambda_B = \frac{x + R(1-x)}{\big(x + R(1-x)\big)y + R(1-y)} \;\leq\; \frac{x + R(1-x)}{xy + R(1-xy)}$$

if we assume $P(e/\neg C \& B) = k$
and $P(e/\neg B) = k$.

for simplicity.

For an ad hoc predictor we take $k = 1$

to get $\Lambda_C = 1 = \Lambda_B$

ie ogives 1 to a enhancement as does evid.

For a non-ad hoc predictor take $k \leq \frac{x}{y}$.

then $\Lambda_C \leq \frac{y}{yx} \leq 1/x$.

Similarly $\Lambda_B \leq 1/y$

so we now get big enhancements leading in our approximation limit to $P(C/e) = 1$ and $P(B/e) = 1$ after confirmation.

this is true regardless of magnitude of x and y.

reward

We turn now to the interesting case of refutation and write

$$P(C/\neg e) = \frac{P(\neg e/e)\,P(C)}{P(\neg e/e)\,P(e) + P(\neg e/\neg C)\,P(\neg e)}$$

where $P(\neg e/e) = P(\neg e/C \& B)\,P(B/C) + P(\neg e/C \& \neg B)\,P(\neg B/C)$

and now consider

$$p(re/cB) = 0$$

$$p(re/_2 e) = 1-k_2, \quad p(re/cB_2\beta) = 1-k_1$$

to give in this case, again assuming $k_1 = k_2 = k$

$$\Lambda_c = \frac{(1-k)(1-y)}{(1-k)(1-y)x + (1-k)(1-x)} = \frac{1-y}{(1-y)x + (1-x)}$$

$$= \frac{1-y}{1-xy}$$

and

$$\Lambda_B = \frac{(1-k)(1-x)}{(1-k)(1-x)y + (1-k)(1-y)} = \frac{1-x}{(1-x)y + (1-y)}$$

$$= \frac{1-x}{1-xy}.$$

~~[To be just the actor could be $k=1$~~

~~Now $\Lambda_c =$~~ Thus formulae for Λ_c and Λ_B are independent of the value of k.

~~Consider $\Lambda_c < 1$ if~~

$1-y < (1-y)x + (1-x)$

or $(1-y)(1-x) < 1-x$ which is always true since $1-y < 1$

~~Hence we have for saturation $\Lambda_c < 1$~~

We note at once that for saturation $\Lambda_c > \Lambda_B$ are always less than unity, as we would expect.

* Footnote

10. The asymmetric effect of rotation in the Bayesian model was reported not to me by Jon Doering who was responsible for the ideas behind the preceding analysis

of successful and unsuccessful productions.

In Fig. 2 ~~we illustrate~~ we illustrate the following sequence — two alternating successes and failures followed by a run of failures when we have assumed as parameter values $x = 0.9$, $y = 0.5$ and $h = 0.1$. The probability P_C for the current and the probability for the current articulation & the protective belt P_B are both shown. After each refutation we take $P_B = 0.5$ again for the new auxiliary hypotheses introduced to account for the anomaly.

We are assuming here the simple situation in which each articulation of the protective belt is triggered by an anomaly. Lakatos (of course stresses cf his [1972] that sophisticated research programmes can progress in an ocean of anomalies which are ignored, the articulation of the protective belt being carried along by the internal momentum of the positive heuristic of the programme. Our ~~interpretation~~ interpretation of Lakatos's methodology, caricature like all rational reconstructions, is a ~~characterisation~~ (compare Lakatos [1972])

but suppose we have x near to 1 and y small

then ~~$\Lambda_e = \frac{1}{x}$~~ for $x=1$, $y=0$ we get

$\Lambda_e = 1$ — hence so is for all values under heat.

But ~~if $x=0$~~ is same heat

$\Omega_B = 0$ so Reflection

acts ~~antisymmetrically~~ asymmetrically on C and B

provided we take x large, y small

as reflecting the mellowing of

upward prejudice.[10] (Formula ~~Λ_e~~ $\times$ mellow

new word explicate

Note in particular we can explicate

rotation without having to

assume ξ & $\hbar$ small.

new word

The upward death of a upward Myanmar

can now be modelled using our

two formulae (10), (12) to allow for the effects

For a successful Rockefeller

(14) $\Lambda_e = \frac{y+\hbar(1-y)}{(y+\hbar(1-y))x+\hbar(1-x)} = \frac{y+\hbar(1-y)}{xy+\hbar(1-xy)}$

and for a rotation

(15) $\Omega_e = \frac{1-y}{(1-y)x+1-x} \quad \frac{1-y}{1-xy}$

Q.E.D.

$x = 0.9$
$y = 0.5$
$h = 0.1$



— P_C
--- P_B

Fig-2.

S = successful prediction
F = failed prediction

newford
Suppose after ~~4th~~ fourth failure in degeneration sequence P_C has fallen to 0.21 and at a stage.)

(P_C must then recover to the value 0.60.)

Every such P_C drop to .06 at 4th, 5th, 6th failure in degeneration sequence ≡ a given limit

Left P_C to 0.26.

(the P_C and P_B curves (marked Q in Fig 2)

Newford
At the earliest point it would become rational for the host cell, to use here an external rational criterion for abandoning a research programme, independent of any competing programme.

new card

At first sight we might suppose that if we have a run of successful predictions followed by a failure then the asymmetry in the effect of failure on P_c and P_B which was large initially would be eliminated as P_c and P_B both now increased to values very close to unity. We shall now show that this rather natural assumption is miscalculated and that in our model the asymmetric effect of failure on P_c and P_B is preserved however many successful predictions precede the failure.

What we shall prove in the following:

After n successful predictions P_c increases from $P_c(0)$ to $P_c(n)$ and P_B from $P_B(0)$ to $P_B(n)$

Let the effect of a failure on P_B, P_c after n successful predictions be denoted by $\Lambda_c(n)$, $\Lambda_B(n)$

Then we find $\frac{\Lambda_B(n)}{\Lambda_c(n)} = \frac{\Lambda_B(0)}{\Lambda_c(0)} = \frac{1-P_c(0)}{1-P_B(0)}$

a result which is independent of n

We have at once the result

$$\frac{\Lambda_B(n)}{\Lambda_c(n)} = \frac{1-P_c(n)}{1-P_B(n)} \qquad \left(\text{see eqn } 15\right)$$

Now use eqn (10) to obtain the result

$$P_c(n+1) = \frac{P_B(n) + P_A\left(1-P_B(n)\right)}{P_c(n)P_B(n) + P_A\left(1-P_c(n)P_B(n)\right)} \times P_c(n)$$

$$P_e(n+1) = \frac{P_B(n)P_C(n) + P_A P_C(n) + P_A P_B(n) P_C(n)}{P_B(n)P_C(n) + P_A - P_A P_B(n) P_C(n)}$$

So $$1 - P_e(n+1) = \frac{P_A(1 - P_e(n))}{P_B(n)P_C(n) + P_A(1 - P_B(n)P_C(n))}$$

Similarly we can show

$$1 - P_B(n+1) = \frac{P_A(1 - P_B(n))}{P_B(n)P_C(n) + P_A(1 - P_B(n)P_C(n))}$$

$$\frac{1 - P_e(n+1)}{1 - P_B(n+1)} = \frac{1 - P_C(n)}{1 - P_B(n)}$$

which applied recursively yields the result

$$\frac{1 - P_C(n)}{1 - P_B(n)} = \frac{1 - P_e(0)}{1 - P_B(0)} = \frac{1 - x}{1 - y}$$

in our old notation

which is independent of n.

Hence we get for rotation after n seconds

$$\frac{\Pi_B(n)}{\Pi_e(n)} = \frac{1 - x}{1 - y} \left(= \frac{1 - P_e(0)}{1 - P_B(0)} \right)$$

In our example $x = .9$, $y = .5$

$\therefore \frac{\Pi_B(n)}{\Pi_e(n)} \doteq \frac{1}{5}$ and ⟶

effect of rotation on B is five times as high

on on C for all values of n,

although of course it is clear that both $P_B(n)$ and $P_C(n)$ both tend rapidly towards 1 with increasing n.

new para

We stress that in this model the corrective effect of refutation on probes for B and for C is clearly accumulated for at all stages by the expected value of P_C being much higher (closer to unity) than we assume for P_B.

6 The Methodological Decision

In the model the effect of refutation on P_C is given a rational (Bayesian) basis.

But other approaches to the methodology of theory evaluation can best be considered as a rather subtle mixture of rational and methodological decisions. We shall now illustrate some of these ideas.

new para Theories faced with an anomaly have various methodological conventions which tell the scientist how to proceed.

So far we have talked about successful predictions, but what happens when a theory makes an unsuccessful prediction, that is to say fixed work at anomaly? There are various methodologies which tell the physicist how to deal with anomalies. For example there is a crude Popperian picture which tells him to reject a theory if it is refuted. But the tenacity of theories is reflected in Lakatos' Methodology of research programmes. To analyze this situation ~~in the context of our Bayesian model again~~ we [have] again wrote T as a conjunction of a hard core C and an auxiliary hypothesis B_1, which belongs to a set $\{B_i\}$ of adjustable hypotheses in the protective belt, adjustable that is in accordance with the positive heuristic of the research programme.

Thus $T = C \& B_1$.

Now suppose the theory T makes a false prediction e_1. Since this is impossible if the theory T were true we must write $P(e_1 / T \& a) = 0$ in equation (3) which leads immediately to $P(T / a \& e_1) = 0$. If we suppose C and B_1 are independent hypotheses for the sake of simplicity, then we can write

$$P(T / a \& e_1) = P(C / a \& e_1) \times P(B_1 / a \& e_1)$$

So we are led to the conclusion $P(C / a \& e_1) = 0$ or $P(B_1 / a \& e_1) = 0$. What alternative we take cannot be decided by the probability calculus alone. It is a matter for methodological [reasoning] with the information [prepared] so far [at our disposal].

decision. Thus the Cabalator modelology can be expressed very simply. We choose to take $P(B_1/z \& e_1) = 0$ and leave $P(c/z \& e_1)$ unchanged in the face of a false prediction e_1. We now invoke a new hypothesis B_2 which credits $c \& B_2$ to explain e_1. This is an ad hoc maneuver for $c \& B_2$ with respect to e_1 so $P(c \& B_2 / z \& e_1)$ can only be enhanced by new successful novel predictions. Suppose e_2 is successfully predicted by $c \& B_2$. Then our previous analysis finds

(14) $$P(c \& B_2 / z \& e_1 \& e_2)$$
$$= \Lambda \times P(c \& B_2 / z \& e_1)$$

where Λ is a suitable enhancement factor.

We can re-write (14) as:

$$P(c/z \& e_1 \& e_2) \times P(B_2/z \& e_1 \& e_2)$$

(15) $$= \Lambda \times P(c/z \& e_1) \propto P(B_2/z \& e_1)$$

But what we actually wished to find was how the credibility of the hard core behaves as the novel programme proposes. To recover this "quotient" we must "distribute" the enhancement ~~factor~~ ratio between the hard core and the protective belt. But again the probability calculus alone cannot

answer this question. Again we have to make a methodological decision. But this decision must be constrained so as not to transgress the rules of the probability calculus. The definition of rates like Poffer's formal account of concepts like reinforcement. We seek a simple formula that satisfies certain conditions. We formulate these conditions for an indefinite number of [next page] In a finite alphabet of attributes related we let $A > B$. stand for two probabilities and we write

$$\Omega\ AB = (\Omega_A \times A) + (\Omega_B \times B)$$

so the enhancement factors Ω ~~Ω_{AB}~~ is elevated between $A > B$ according to the factors Ω_A and Ω_B which will be functions of Ω, as well as each depending on $A > B$.

Regarding $\Omega_A > \Omega_B$ as functions of Ω we list the following ~~natural~~ restrictions, assuming $A > B$ without loss of generality

(1) $\Omega_A(1) = \Omega_B(1) = 1$

(2) $\Omega_A\left(\frac{1}{AB}\right) = 1/A$, $\Omega_B\left(\frac{1}{AB}\right) = 1/B$

(3) $\Omega_A(\Omega)$ and $\Omega_B(\Omega)$ are monotonically increasing functions of Ω on the range $\left(1, 1/AB\right)$

(4.) $\Omega_A \times A > \Omega_B \times B$ ~~if $A > B$~~ for all Ω in range $(1, 1/AB)$

(5) $\Omega_A + \Omega_B = \Omega$.

(6) Writing $\Omega_A(\Omega) = f(A, B, \Omega)$

$$f(A, B, \Omega'\Omega'') = f(A, B, \Omega') \times f\left[f(A, B, \Omega')A, \frac{\Omega}{f(A, B, \Omega')}B, \Omega''\right]$$

must satisfy conditions on $\Omega_{B/A}(\Omega)$

Condition (2) arises from setting that the maximum value of Ω is $1/AB$ since we require that Ω_{AB} is a probability, the maximum value for Ω_A is $1/A$ since $\Omega_A A$ is to be a probability, and similarly for Ω_B.

Condition (4) means that enforcement now leads to a "clear-cut" case of the assessment of A and B.

Condition (6) means that our rule for evidencing at Ω_A can be built up consistently in stages.

Conditions (1), (3) and (5) require no comment.

It is perhaps surprising that a simple formula for Ω_A can be obtained satisfying all these conditions. viz

(18) $$\Omega_A = \frac{\Omega(1-AB)}{1-A+A(1-B)\Omega}.$$

Ω_B is then just twice as on Ω/Ω_A.

(17) i.e. $$\Omega_B = \frac{1-A+A(1-B)\Omega}{1-AB}$$

For $A \angle B$ we interchange the formulae for Ω_A, Ω_B at the same time exchanging A and B in these formulae

Exercise 17.

Exercise

* 9 am grateful to Jon Dollery for drawing my attention to following result concerning solution for ~~Exercise~~ exercises 1 to 3.

$$\rho_A = \frac{B + R(1-B)}{AB + R(1-AB)}$$

where R is the unique root in the range $0 < R < 1$ of the following quadratic equation

$$R^2\{(1-A)(1-B) - (1-AB)^2\rho\}$$
$$+ R\{A + B - 2AB - 2AB(1-AB)\rho\}$$
$$+ AB(1-AB\rho) = 0.$$

We leave the derivation of the result as a challenge to the interested reader.

It is conjectured that (16) > (17) are the only simple solutions satisfying the conditions (1) to (6)

The only other property we might like to impose is

(7) $\lambda_A = \lambda_B = \sqrt{\lambda}$ for $A = B$.

Our solution (16) > (17) does not satisfy (7) condition

and it may even be the case that conditions 1 to 7 are incompatible.

There appears to be no simple solution that satisfies all 7 conditions (1) to (7). Note *12

Hence we are only interested in a model which will keep the general features of different realizations (16 and 17) we shall adopt our solutions (16) > (17)

Another feature we might want to build into our model is a memory (Footnote 13) of the past. Not to say after a number of unsuccessful iterations we do not throw to the probability of the Pard are entirely discharged but "depress" it at the m^{th} iteration by a memory factor $F(m)$ which began to act at the critical value m_0.

We can now put all these features together to obtain a model for the following very general problem

13. I am grateful to Noetta Kintop for suggesting this.

new para

Suppose a research programme makes n_m successful predictions following the $(m-1)^{th}$ false prediction by suitable articulation of its positive heuristic

new para

We denote by $P_c(n_1, n_2 \ldots n_m)$ the posterior probability for the hard core c after this sequence of successes and failures

Then we can write the following recursion for P_c

$$P_c(n_1, n_2, \ldots n_m) = P_c'(n_1 \ldots n_{m-1}) \times \textcircled{H}\left(\sqrt{x}, P_c'(n_1 \ldots n_{m-1}), \Lambda^{(n_m)}\right)$$

where $P_c'(n_1 \ldots n_{m-1}) = F(m-1) \times P_c(n_1 \ldots n_{m-1})$

and $\Lambda^{(n_m)} = \underline{\Phi}\left(\varepsilon, n_m, \sqrt{x}\, P_c'(n_1 \ldots n_{m-1})\right)$

$\Lambda^{(n_1)} = \Phi(\varepsilon, n_1, x)$

with $P_c(n_1) = \sqrt{x}\, \textcircled{H}\left(\sqrt{x}, \sqrt{x}, \Lambda^{(n_1)}\right)$

and $\textcircled{H}(A, B, c) = \dfrac{c(1-AB)}{1-B+B(1-A)c}$

$\underline{\Phi}(A, B, c) = \dfrac{1}{c(1-AB)+AB}$

$F(m) = \dfrac{1}{1+\exp(m-m_0)^{\frac{m}{\sigma}}}$

In these formulae as before $x = p(C \& B_i \mid \bar{b})$ and

$\varepsilon = p(e / r(C \& B_i))$.

We have assumed that initially equal probabilities $\sqrt{x}$ are attached to C and B_i, and that each successive articulation of the protective belt starts again with probability $\sqrt{x}$ ~~although what happens to Pc after refutation will depend on our methodology~~.

new para We have chosen a simple form for $F(n)$ with suitable initial properties.

new para Of course if the number of alternative hypotheses in the protective belt that can be invoked to deal with anomalies is a finite number N say, then taking the pure Lakatos model without a memory factor, Pc will drop immediately to zero after $N+1$ anomalies have displaced zero no further articulation of the protective belt is possible.

new para Taking the parameter values $N=6$, $n_0 = 3$ $\sqrt{x} = 0.01$ $\varepsilon = .1$ and assuming for purposes of illustration the case where successful and unsuccessful predictions alternate, the account derived from our model is shown in fig 3 below, once for pure Lakatos (no memory factor) and pure Popper (refutation takes us back to square one, i.e. Pc falls to $\sqrt{x}$ after each refutation) are also given for comparison.

new para Certainly the memory models looks the

most realistic although we have chosen a very simple type of money — a more sophisticated version would take due account of small or vanishing values of N_m for as m increases, corresponding to the informal notion of legendary in the research programme.

At all events we have succeeded in creating a coherent theory of demand for Boffinair models for the life and death of a research programme



Fig. 3.

$N = 6$
$m_0 = 3$
$\sqrt{x} = .01$
$\varepsilon = .1$

I pure Lakatos : refutation leaves P_c unchanged

II Lakatos plus memory ; refutation cuts P_c by a memory factor ($\sqrt{x}$ Hm).

III Pure Popper. : refutation takes P_c back to $\sqrt{x}$.

new para

Let us now briefly compare our new model with the procedure presented in Section 5. In the first place the new model assumes that Rand acts independently on the people Bayesian model. Of course the first model suffers more significantly from more complicated conditional probabilities such as $P(re/BBnC)$ etc. Given the second model assumes we know only the value of $P(re/nBvnC)$ or our input information.

Naturally our methodological decisions in the second model could be re-examined. Conditional on the assignments attained in the first model - (~~this~~ in fact ~~is~~ true during ~~around~~ at his solution ~~I~~ our conclusion ~~to~~ ~~Re~~ for the second model) But the states of these assignments is now quite different. For the first model we denied the effect of refutation from these variables in the second model we effectively constrain these assignments ~~from~~ by the decisions we chose to ~~take~~ make about the effects of refutation.

Finally we note that in the second model we can explain the progress ~~in~~ of rational ~~even~~ allowing P_B and P_C initially to have the same low value, irrespective of success & of first

Model depends critically on taking steps as required.

At all events we believe that we have successfully distinguished the possible roles of Bayesian & semi-Bayesian models in accounting for the life and death of a research programme.

References Lakatos I [1970]:

Redhead, M.L.G. [1974]: 'On Neyman's Paradox and the Theory of Statistical Tests' Brit. J. Phil. Sci, 25 pp. 265-271.

Salmon, W.C. [1967]: The Foundations of Scientific Inference.

Schaffner, K.F. [1974] 'Einstein versus Lorentz: Research Programmes and the Logic of Comparative Theory Evaluation'. Brit. J. Phil Sci 25 pp. 45-78

Zahar, E.G. [1973]: 'Why did Einstein's Programme Supersede Lorentz's?' Brit. J. Phil. Sci, 24 pp. 95-123 and 223-62.

Laudrap B.E., Peterman, A. and de Rhoplee, F. [1972]: 'Comparison between Theory, Experiment in quantum Electrodynamics' Physics Reports, 3C (1972), 195-259.

Rich, A. and Wesley J.C. [1972] 'The current status of the lepton g factors' Rev Mod. Phys. 44, (1972) pp. 250-283.. (Reviews of Modern Physics)

Lakatos I [1970]: 'Falsification and the Methodology of Scientific Research Programmes', in I. Lakatos and A. Musgrave (eds): Criticism and the Growth of Knowledge, pp 91-195.

[Please write Brit. J. Phil Sci in full (Marie-Louise Thas:- The British Journal for the Philosophy of Science- in references for Redhead [1974], Schaffner [1974] and Zahar [1973]]